



Process parameters optimization and molecular studies on bacteriocin-producing lactic acid bacteria from 'Kati'

Aarinade Adejumo Omotayo, Oladipo Oladiti Olaniyi*, Bamidele Juliet Akinyele

Department of Microbiology, Federal University of Technology Akure, Nigeria.

ARTICLE INFO

Article history:

Received 17 Oct. 2018
Received in revised form
21 Feb. 2019
Accepted 02 Mar. 2019

Keywords:

Crude bacteriocins;
Lactic acid bacteria;
Bacterial pathogens;
'Kati';
Antibacterial activity

ABSTRACT

This work described the isolation and identification of bacteriocin-producing lactic acid bacteria (LAB) from 'Kati' (a sorghum based fermented food), and to evaluate the antibacterial effect of bacteriocin on selected pathogenic bacteria. The identities of the isolates were revealed to be as *Lactobacillus plantarum*, *L. brevis*, *L. fermentum* KAT1, *L. fermentum* KAT2 and *Lactococcus lactis* using 16S rRNA gene sequence analysis. Out of 28 LAB, five were found to inhibit selected pathogenic bacteria namely; *Escherichia coli*, *Staphylococcus aureus*, *Salmonella typhimurium* and *Bacillus cereus*. The unpurified bacteriocins produced by the isolated LAB were characterized with respect to the effect of temperature, pH and surfactant. The test isolates showed activities of 6400, 6400, 3200 and 1600 AU/ml respectively. Crude bacteriocin from *L. brevis* and *L. lactis* were the most heat stable at 121°C for 60 min. Bacteriocins from *L. plantarum*, *L. lactis* and *L. fermentum* KAT1 showed the highest antibacterial activity and stability at pH 2.0 to 6.0. Exposure to Tween 20 increased the bacteriocin activity of the LAB isolates except for *L. fermentum* KAT2 where loss of activity occurred. The findings from this study suggest that bacteriocinogenic LAB present in 'kati' have potentials to inhibit pathogenic/spoilage microorganisms in foods.

Citation: Omotayo AA, Olaniyi OO, Akinyel BJ. **Process parameters optimization and molecular studies on bacteriocin-producing lactic acid bacteria from 'Kati'**. J Food Safe & Hyg 2019; 5(1): 10-18

1. Introduction

The roles of lactic acid bacteria (LAB) are vital in the production of numerous fermented foods globally. Arrays of strains are frequently utilized as inocula in the fermentation of milk, cereals meat and vegetable (1). During the course of fermentation, LAB exerts desired organoleptic qualities on substrates being fermented and extends their shelf life by secreting antibacterial metabolites against food spoilage microorganisms (2). The antibacterial effect of LAB has been exploited by people for hundreds of years without any adverse effects on consumers. It has improved the nutritional profile, shelf life and safety of fermented foods. The preservation of fermented foods by LAB is connected to the secretion of certain antimicrobial

substances which includes bacteriocins, organic acids, hydrogen peroxide, reuterin and diacetyls. Among the various antimicrobial compounds secreted by LAB, bacteriocins take the lead in terms of antagonism against spoilage organisms commonly implicated in foods (3). Examples of lactic acid bacteria are: *Lactobacillus* spp, *Leuconostoc* spp, *Lactococcus* spp, *Streptococcus* spp, *Enterococcus* spp, *Pediococcus* spp, *Weissella* spp, e.t.c. An arrays of bacteriocins from LAB that have been identified and characterized include Nisin, Diplococcin, Acidophilin, Bulgarican, Helveticins, Lactacins, and Plantaricins. Out of these bacteriocins, Nisin produced by *Lactococcus lactis* sp. *Lactis* has been commercialized and extensively used (4).

LAB have a long historical application in food production due to its profound effects on organoleptic

*Corresponding author. Tel.: +2348068054636
E-mail address: microladit@gmail.com

and nutritional properties (5). They acidified raw materials via the production of organic acids, mainly lactic acid. In addition to the production of organic acids, acetic acid, aromatic substances, ethanol, bacteriocins, exopolysaccharides and relevance extracellular enzymes are produced by LAB. Traditionally, food fermentation process by LAB is based on spontaneous. However, in an industrial setting, food matrix is deliberately inoculated with LAB as starter cultures under controlled cultural parameters. The recent use of high profile novel starter cultures in food fermentation process is being explored for the purpose of improved functionalities and standardization of the end products (6). Also, certain strains of LAB have been noted to possess some desired health-promoting qualities; one of such qualities of these probiotics is the ability to either eliminate or inhibit the growth of some bacterial pathogens in the gastrointestinal tract. Few of the pathogens that have been successfully combated by potential probiotics are *Helicobacter pylori*, *Escherichia coli*, and *Salmonella* (4).

In recent years, extensive research has targeted bacteriocin-producing strains of LAB with improved properties due to their food preserving potential connected with the antimicrobial nature of bacteriocins (7). Bacteriocin-producing LAB with a broad spectrum are of special interest in food industry because these bacteria are generally regarded as safe (GRAS). They occur naturally in fermented foods in a wide range of applications in food industry. Nowadays, there is a growing concern regarding the replacement of the use of chemical preservatives in food. LAB have been in use for a while for the fermentative preservation of several food products from dairy product to meat and vegetables. They can make antimicrobial substances (e.g. organic acids, hydrogen peroxide and bacteriocins) which can instigate the growth of the potential harmful and/or spoilage microorganisms. Since vegetables are sources of several vitamins, dietary fibers and minerals, it is desirable that vegetable based food products preserve their biological values during fermentation. Besides the traditional role of LAB in food preservation, they exert beneficial effects to the gut. Manufacturing food products with long shelf life and viable bacteria having confirmed health promoting result can contribute on a large scale to the human well-being in the near future.

Recent concerns about emerging microbial populations with antibiotic resistance and undesirable toxic properties of several bioactive peptides stimulated attention to the bacteriocins as promising natural antimicrobials in combating pathogens and

spoilage microorganisms. To accomplish this task, a precise understanding about the nature of these molecules is required (8). For now, the only commercially available and marketed bacteriocin is nisin and it is more active against Gram-positive spore-forming bacteria (3). Other bacteriocins of *Lactobacillus* spp. have been reported to be active against closely related species of mesophilic *Lactobacillus* and therefore regarded as potential natural food preservatives (9). Scanty information and limited use of LAB with antibacterial properties inform this present study. This study is therefore designed to screen bacteriocin-producing LAB isolated from 'kati': a sorghum based fermented food for bacteriocin production and evaluate their antibacterial effect on food spoilage bacteria.

2. Materials and methods

2.1. Sample collection

'Kati' samples were purchased from four different locations namely Ikare, Akungba, Oka-Akoko and Okeagbe in Akoko area of Ondo state, Nigeria. The sorghum sample (control) was obtained from Oja-Oba Market in Akure, Ondo State, Nigeria making 15 samples in all. The samples were deposited in an aseptic polythene bags and then conveyed to Microbiology Laboratory, FUTA for further microbiological and chemical analysis.

2.2. Preparation and production of 'kati'

Sorghum grains were sorted, soaked in distilled water for three days and then wet-milled. After milling, the milled dough was fermented for 12 h and were wrapped with cocoa (*Theobroma cacao*) leaves and then cooked for 20-30 min. The cooked dough was moulded vigorously in between the palms and then wrapped in fresh cocoa leaves and cooked again for 30-45 min. The cooked dough was cooled and kept in a sterile polyethylene bag. The stored cooked dough is now 'kati'.

2.3. Bacterial strains and culture

One gram of each 'kati' sample (raw, fermented and cooked sample) was weighed and dispensed separately into test tubes each containing 9 ml sterile distilled water and serially diluted up to 10^{-10} dilution factor. From the diluents, 0.1 ml aliquot from the dilution factor 10^{-3} and 10^{-5} of each sample was dispensed into different sterile Petri dishes containing deMan rogosa agar (MRS), allowed to solidify and incubated anaerobically at 30°C for 48 h. The cultures

were purified by repetitive streaking.

2.4. Production of crude bacteriocin samples

Lactobacillus species were grown in 400 ml MRS agar broth for 72 h at 30°C anaerobically in triplicate. For extraction of bacteriocin, a cell-free solution was centrifuged (4000 rpm for 40 min at 4°C) and was adjusted to pH 7.0 by 1 M NaOH to eliminate organic acids. The supernatant was thereafter filtered using 0.2 µm pore-size membrane filter. Inhibitory activity from hydrogen peroxide was removed by the inclusion of 2 mg/ml catalase (10).

2.5. Determination of bacteriocin activity

Agar well diffusion method was used as illustrated by Takahiro et al. (11). 5µl crude bacteriocin from the LAB strains from each dilution were placed in wells in plates implanted with the indicator/test strain; incubated aerobically at 37°C for 24 h and the zone of inhibition diameters were recorded (12).

2.6. Determination of bacteriocin titre

The quantification of bacteriocin titre was performed by two fold serial dilutions of bacteriocin in saline solution and aliquots of 50 µl from each dilution were put in wells in plates incorporated with the indicator strains. The cultures were then incubated for 18-24 h at 37°C under aerobic condition and the zone of clearance (2 mm and above) which indicate inhibition of indicator organisms around the wells were observed and measured. The antibacterial activity of the bacteriocin was defined as the reciprocal of the highest dilution showing inhibition of the indicator lawn and was expressed in activity units per ml (AU ml⁻¹) (13).

2.7. Molecular identification of bacterial isolates via 16S rRNA Gene Sequencing method

The identities of the pure bacterial cultures were authenticated by molecular tool. The DNA from pure bacterial cultures was isolated by Fast DNA bacterial isolation Kit in accordance with the manufacturer's instruction. The 16S rRNA gene in the isolated bacterial DNA was amplified by Polymerase Chain Reaction (PCR) using two universal 16S rRNA primers (14). The 50 µl PCR mixtures contained 1 µl DNA template, 25 µl 2 × ExTaq PCR (Takara Bio INC, Shiga Japan), 1 µl of each primer ((Bioray biotechnology, Xiamen, China) and 22 µl double-distilled water. The PCR

procedure adopted was as follows: primary denaturation for 5 min at 95°C; 30 cycles of denaturation at 94°C for 30 s; annealing at 58°C for 30 s, and extension at 72°C for 1 min; and an additional reaction for 10 min at 72°C. The quality of PCR products were inspected by agarose gel electrophoresis after which they were submitted Bioray Biotechnology (Xiamen, China) for sequencing. The 16S rRNA gene sequence obtained was distinguished with other 16S rRNA gene sequences already deposited in GenBank via BLASTN program (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) and aligned with similar sequences using CLUSTX program (15).

2.8. Characterization of bacteriocin

In order to ascertain the optimum temperature and pH, and the effect of surfactant on crude bacteriocin activity, the samples were characterized by subjecting them to varied conditions. Their stability at different conditions was also evaluated.

2.8.1. Heat resistance

Crude bacteriocin was subjected to different incubation temperatures at 40, 60, 80, 100 and 121°C. At 0, 30, 60 or 90 min of incubation, aliquots were withdrawn and bacteriocin activity was assayed (16).

2.8.2. pH sensitivity

Crude bacteriocin were regulated to pH 2, 4, 6, 8, 10, and 12 with HCl and NaOH, incubated for 4 h at room temperature and assayed accordingly (16).

2.8.3. Effect of surfactant on bacteriocin activity

The effect of surfactant on bacteriocin activity was investigated by integrating non-ionic surfactant (tween 20). The surfactant was included into crude bacteriocin at a concentration of 0.1 ml of surfactant ml⁻¹ of bacteriocin solutions. This preparation was incubated at 30°C for 60 min and assayed for bacteriocin activity against the food spoilage bacteria (17).

3. Results

3.1. Antibacterial activity of crude bacteriocin from isolated LAB against some pathogenic bacteria

The selected LAB strains from 'kati' produced bacteriocin which revealed antibacterial activity against two Gram-positive and two Gram-negative bacterial strains (Fig 1). The crude bacteriocin from

selected LAB showed varied inhibition against all the indicator organisms. The crude bacteriocin from *L. lactis* showed the highest inhibitory effect against *Bacillus cereus*, *Staphylococcus aureus* and *Salmonella typhimurium*, while the crude bacteriocin from *L. plantarum* had the highest antibacterial activity against *Escherichia coli*. *E. coli* was noted to have the highest susceptibility out of the four indicator organisms to the crude bacteriocin from *L. plantarum*, *L. lactis* and *L. brevis*.




Figure 1. Antibacterial activity of crude bacteriocin from isolated LAB against some pathogenic bacteria

3.2. Genomic characterization of the LAB isolates

Molecular identities of the LAB via 16S rRNA Sequence analysis are presented in Table 1. The identities of these strains were revealed by comparing the gene sequences obtained with those available in GenBank. The LAB were identified as *Lactobacillus plantarum*, *L. lactis*, *L. brevis*, *L. fermentum* KAT1 and *L. fermentum* KAT2.

Table 1. Molecular identities of the LAB

Cultural and biochemical Identification	Gene sequence identification	Max identity (%)	Accession number
<i>Lactobacillus</i> spp	<i>Lactobacillus plantarum</i>	97%	AY096004.1
<i>Lactobacillus</i> spp	<i>Lactococcus lactis</i> L.	97%	CP020604.1
<i>Lactobacillus</i> spp	<i>brevis</i>	99%	CP012110.1
<i>Lactobacillus</i> spp	<i>L. fermentum</i> KAT1	95%	CP011536.1
<i>Lactobacillus</i> spp	<i>L. fermentum</i> KAT2	96%	CP002033.1

3.3. Effect of temperature on antibacterial activity of bacteriocin from isolated LAB

The effect of temperature on the bacteriocin activity was evaluated by incubating the crude bacteriocin (at 40, 60, 80, 100 and 121°C) for various periods (Fig 2-6). The inhibitory activity of the bacteriocin from *L.*

plantarum incubated at different temperatures for different time interval is shown in Fig 2. At 40°C for 10, 30 and 60 min of incubation, the inhibitory activity of the bacteriocin from *L. plantarum* against *B. cereus* was relatively stable and at 90 min the antibacterial spectrum reduced from approximately 6400 AU/ml to 3200 AU/ml. The bacteriocin activity of *L. plantarum* was heat stable at 60 and 80°C from 10 to 30 min and at 60 to 90 min bacteriocin activity was reduced. At 100°C, the bacteriocin activity was stable at 10 min, but reduced at 30 to 60 min, the activity was further reduced at 90 min. At 121°C, *L. plantarum* showed activity of 3200, 1600, 400 AU/ml at 10, 30 and 60 min respectively. While at a temperature of 121°C for 90 min, there was loss of activity.




Figure 2. Effect of temperature on bacteriocin activity from *L. plantarum* at different time interval

The bacteriocin activity of *L. lactis* shown in Fig 3. At 40°C, the bacteriocin activity was stable at different time interval of incubation, while at 60°C, bacteriocin activity was stable from 10 to 60 min but reduced at 90 min. At 80 and 100°C, the bacteriocin activity resulted in 6400 AU/ml for 10 min and was reduced to 3200 and 1600 AU/ml from 30 to 90 min. At 121°C, the bacteriocin activity was 3200, 3200, 1600 and 400 AU/ml at 10, 30, 60 and 90 min respectively.

L. brevis showed a significant bacteriocin activity at different temperature range and time interval of incubation (Fig 4), it was the only test isolate that exhibited highest activity at 121°C at all the time intervals which made it the most heat stable out of all the five isolated tests. The bacteriocin activity shown by *L. brevis* ranged from 6400 AU/ml to 3200 AU/ml at 40, 60, 80 and 100°C for 10, 30, 60 and 90 min. While at 121°C, it was stable at 10 min, but reduced to 3200 AU/ml at 30 min and reduced further

to 1600 AU/ml from 60 to 90 min.




Figure 3. Effect of temperature on bacteriocin activity from *L. lactis* at different time interval




Figure 4. Effect of temperature on bacteriocin activity from *L. brevis* at different time interval

L. fermentum KAT1 demonstrated considerable bacteriocin activity at temperature of 40, 60 and 80°C but mostly at 40°C at different time interval which is shown in figure 5. However, the bacteriocin activity ranged from 6400 to 400 AU/ml. At 100°C, there was bacteriocin activity from 10 to 30 min but no activity was shown from 60 to 90 min. Furthermore, there was complete loss of activity at 121°C at all the time interval.

The bacteriocin of *L. fermentum* KAT2 is shown in figure 6. At 40°C, the antibacterial activity was stable from 10 to 60 min, while at 90 min, it reduced 6400 to 3200 AU/ml. At 60 and 80°C, the bacteriocin activity was stable from 10 to 30 min, but reduced from 60 to




Figure 5. Effect of temperature on bacteriocin activity from *L. fermentum* KAT1 at different time interval

90 min. At 100°C, the activity was 6400, 3200, 3200 and 1600 AU/ml at 10, 30, 60, and 90 min, respectively. There was considerable bacteriocin activity from 10 to 60 min, while at 90 min, there was loss of activity.




Figure 6. Effect of temperature on bacteriocin activity from *L. fermentum* KAT2 at different time interval

3.4. Effect of pH on antibacterial activity of bacteriocin from isolated LAB

Supernatant samples taken after centrifugation were adjusted to different pH and were shown to be effective against the test organism *B. cereus*. The impact of pH on bacteriocin activity is shown in Fig 7. Bacteriocins exhibited antibacterial efficacy and were stable in an acidic pH ranged from 2 to 6, while at pH 8 to 12, the antibacterial activity was minimal and the stability was reduced. At pH 4, the bacteriocin activity of all the LAB isolates was stable. The highest bacteriocin activity was demonstrated by *L. lactis*. The least bacteriocin activity was shown by *L. brevis*. The

bacteriocin activity shown ranged from 6400AU/ml to 400AU/ml.




Figure 7. Effect of pH on bacteriocin activity from the lactic acid bacteria

3.5. Effect of surfactants on antibacterial activity of bacteriocin from isolated LAB

Exposure to surfactants (0.1% Tween 20) gave rise to an increase in the bacteriocin activity rather than decrease in activity. Fig 8 reveals the effect of Tween 20 on bacteriocin activity of the isolated lactic acid bacteria. Significant bacteriocin activity was exhibited by the *L. plantarum* and *L. lactis*, with an activity of 25600 AU/ml. A moderate activity was shown by *L. brevis* and *L. fermentum* KAT1 while *L. fermentum* KAT2 showed loss of activity.




Figure 8. Effect of surfactant on bacteriocin activity of the isolated lactic acid bacteria

4. Discussion

The generation of a bacteriocin with a vast spectrum of antibacterial efficacy against similar or

closely related bacterial strains can be an essential attribute for the selection of LAB as starter cultures and is of significance in the fermentation of cereals and legumes. The adoption of bacteriocins from LAB has been reported to enhance the nutritional and organoleptic quality of fermented foods and also reduce the processing cost. At present, certain strains of *Lactococcus lactis* and *Pediococcus* spp. are known to be producers of Nisin and Pediocin PA1 respectively. Nisin being the most extensively applied bacteriocin, it has been considered as GRAS (generally recognized as safe) in the United States for food (18). In the last few years, high premium has been placed on bacteriocins from LAB as a promising candidate in place of chemical preservatives. They are regarded as effective bio-preservatives agents and their application in food is presently the subject of thorough research.

The current investigation accentuates the isolation and identification of LAB strains from 'kati', production and characterization of bacteriocins produced by them. In this study, five wide spectrum bacteriocin-producing LAB strains were encountered and identified as *L. brevis*, *L. plantarum*, *L. lactis*, *L. fermentum* KAT1 and *L. fermentum* KAT2. Most of the LAB strains isolated and identified in this work had been previously isolated from some other fermented cereals with *L. plantarum* as the predominant organism. Ejikeme and Ijeoma (19) isolated *L. plantarum*, *L. cellobiosus*, *L. pentosus*, *Leuconostoc mesenteroides* and *Pediococcus pentosaceus* from *ogi*, a cereal-based fermented product, while Julius et al. (20) isolated *L. plantarum* and *L. mesenteroides* from *maasai*, a fermented maize dough. In another study by Fowoyo and Ogunbanwo (21), an array of LAB to include *L. fermentum*, *L. plantarum*, *P. acidilactici*, *L. lactis* and *L. mesenteroides* were identified from the various fermenting stages of white maize obtained from the South-western region of Nigeria for *massa* production, a traditional Nigerian snack made from fermented maize grains. The production of organic acids in the fermentation medium accounts for the presence of acid tolerant LAB to the detriment of the competing organisms (22).

The antibacterial spectrum of bacteriocin from LAB isolated from 'kati' against selected bacterial pathogens was evaluated using *in vitro* technique. Out of the entire 28 LABs, only 5 strains showed antagonistic effect against the indicator organisms, after which molecular identification was carried out on Bacteriocin produced by the *Lactobacillus* spp. produced obvious and clear zones of inhibition against the selected pathogenic/spoilage bacteria. Bacteriocins from the

isolated LAB have a few striking qualities that validate this study. The most remarkable is that none of these bacteriocins is limited by the immensely narrow antibacterial spectrum reported for some bacteriocins of some LAB, for example Lactococcin A (23) and Lactacin B (24). The highest degree of inhibition was exhibited by *L. lactis* and *L. plantarum*. Earlier reports (25) have demonstrated that some bacteriocins produced by gram-positive bacteria have a better antibacterial activity.

The inhibitory activity of the bacteriocin from the LAB was fairly stable at different temperatures but decreased with increase in the incubation time. The antibacterial activity of bacteriocins was fairly stable in all the time intervals (minutes) at 40, 60, 80 and 100°C, but was completely lost after incubating at 121°C for *L. fermentum* KAT1. The bacteriocin from *L. brevis* was most heat stable as there was no appreciable decline in activity after heating at 121°C for 60 min whereas, there was notable decline of activity following heat treatment at 121°C for all time intervals for both *L. fermentum* KAT1 and *L. fermentum* KAT2. Although heat stability of antibacterial activity of bacteriocin produced by *Lactobacillus* spp has been well documented (26). The stability in the antibacterial activity of bacteriocin produced by *L. brevis* subjected to heat treatment at 121°C for 60 min is exceptional. *L. plantarum* as well as *L. lactis* showed significant heat stability. Heat stability is crucial for any bacteriocin to be used as a food preservative, because many mechanisms of food production require a heating process. Heat resistance is an important quality of several bacteriocins secreted by LAB and can differ effectively ranged from 60°C or 100°C for 30 min and above at 121°C for 15-20 min (9). Different bacteriocins produced by LAB, especially the class I and class II, are portrayed as small hydrophobic proteins comprising small tertiary structure, which clarifies their heat stability. Other determinants contributing to heat stability of the bacteriocin of LAB are stable cross-linkages, a high glycine content and occurrence of strongly hydrophobic regions (9). Studies have shown and indicated that cold temperature is the most suitable for the preservation of bacteriocins for an effective application in the food industry. Therefore, the unique antibacterial attributes of *L. brevis*, *L. plantarum*, *L. lactis*, *L. fermentum* KAT1 and *L. fermentum* KAT2, can definitely have a great effect on their application as starter cultures for the production of fermented foods, with a perspective to bettering the cleanliness and safety of the foods produced.

Bacteriocins from the lactic acid bacteria exhibited antibacterial activity against test organisms and were stable in the pH ranged between 2 and 6, while its activity declined in alkaline regions (pH 8 to 12). The action of bacteriocin from the isolated LAB was demonstrated to be pH dependent. Bacteriocins produced by *L. plantarum*, *L. lactis* and *L. fermentum* KAT1 showed the greatest activity and stability at pH 2-6. Remarkable acid stability was earlier exhibited by plantaricin, bulgarican and lactobulgarican (24). The highest antibacterial activities of bulgarican and lactobulgarican from *L. bulgaricus* were reported to be stabled at pH 2.2 and 4.0 respectively against arrays of food spoilage and pathogenic bacteria. Bacteriocins vary tremendously with respect to their stability in different pH and temperature. Many of the bacteriocins produced by LAB are only stable at acid and neutral pH (9) and are inactivated even at a pH above 8.0 (e.g. nisin, lactostrepcins, pediocin AcH, leucocin A-UAL 187). The bacteriocin produced by *L. lactis* was stable at alkaline pH values which makes them an appealing subject in food supplies. A relevant detriment while using bacteriocins is in the likely decline of their longstanding efficiency attributed to pH changes, specifically in fermented foods. In this regard, bacteriocin activity of the isolated LAB were not altered by changing in pH; activity levels were still high at acidic and basic pH values.

Inhibitory activity was not lost following treatment with surfactant (Tween 20). Exposure of the bacteriocin in this study to surfactants (Tween 20) resulted in significant increase in bacteriocin activity. *L. plantarum* and *L. lactis* showed the highest activity after the addition of surfactants to the bacteriocin while *L. fermentum* KAT2 showed no activity with addition of the surfactants (Tween 20). It was reported by Malheiros et al. (27) that the supplementation of the bacteriocin production medium with Tween 20 resulted in highest bacteriocin yield by *L. sakei*. In a similar study by Castro et al. (28), the addition of Tween 20 to MRS broth enhanced *L. curvatus* ACU-1 bacteriocin activity. Tween 20 is a non-ionic surfactant agent capable of increasing bacteriocin production due to its effect on membrane fluidity and stimulation of the secretion of proteins (29). Inhibitory activity of bacteriocin might be due to the effect of surfactant in the permeability of the cell membrane of the test isolates (9).

5. Conclusion

16S rRNA sequence analysis showed that *kati* contains diversity of LAB. The crude bacteriocin from the LAB strains expressed antibacterial activity against all the indicator organisms. The antibacterial activities of the bacteriocins were fairly stable over a wide range of temperature and pH. Hence, they could be exploited for various industrial purposes as bio-preservatives since various criteria useful for applications are met. It is therefore suggested that these bacteriocins could be employed as a productive control for spoilage and pathogenic microorganisms as they were able to show antimicrobial activity.

Conflict of interest

The authors declare they have no conflict of interest.

Acknowledgments

The technical supports from the Department of Microbiology, Federal University of Technology Akure, Nigeria is highly appreciated.

References

1. Wood BJ, Holzapfel WH. The Genera of lactic acid bacteria. Vol. 2, Blackie Academic and Professional, London. 2010.
2. Buddle BB, Hornbaeck T, Jacobson T, et al. *Leuconostoc carnosum* 4010 has the potential for use as a protective culture for vacuum-packed meats: Culture isolation, bacteriocin identification, and meat application experiments. *Int J Food Microbiol* 2003; 83: 171-84.
3. Deegan LH, Cotter PD, Hill C, et al. Bacteriocins: Biological tools for biopreservation and shelf-life extension. *Int Dairy J* 2013; 16: 1058-71.
4. Leroy F, De Vuyst L. Lactic acid bacteria as functional starter cultures for the food fermentation industry. *Trends Food Sci & Technol* 2010; 15: 67-78.
5. Chelule PK, Mokoena MP, Gqaleni N. Advantages of traditional lactic acid bacteria fermentation of food in Africa. *Curr Res Technol Education Tropics in Appl Microbiol & Microbial Biotechnol* 2010; 1160-67.
6. Pedreno M, Gandia F, Caballero N, et al. Characterization of the antioxidant activity of betalains from *Beta Vulgaris* L. roots Neo diet-cost. *Muscia Spain*. 2013; 3: 916-8.
7. Savadago A, Quattara CT, Bassole IH, et al. Bacteriocins and lactic acid bacteria - A mini-review. *Afri J Biotechnol* 2012; 5: 678-83.
8. Asahara T, Nomoto K, Watanuki M, et al. Antimicrobial activity of intraurethrally administered probiotic *Lactobacillus casei* in a marine model of *Escherichia coli* urinary tract infection. *Antimicrob Agents & Chemo* 2011; 45: 1751-60.
9. De Vuyst L, Vandamme EJ. Antimicrobial potential of lactic acid bacteria. In: *Bacteriocins of lactic acid bacteria*. Springer US 2010; 91-142.
10. Daba H, Pandian S, Gosselin JF, et al. Detection and activity of bacteriocin produced by *Leuconostoc mesenteroides*. *Appl Environ Microbiol* 2000; 57: 3450-55.
11. Takahiro KA, Sheldon BW, Klapes NA, et al. Nisin treatment for inactivation of *Salmonella* species and other Gram negative bacteria. *Appl Environ Microbiol* 1991; 57: 3613-15.
12. Rammelsberg M and Radler F. Antibacterial polypeptides of *Lactobacillus* species. *J Appl Microbiol* 2002; 69: 177-84.
13. Graciela H, Lacroix C, Huang J, et al. Simple method of purification and sequencing of a bacteriocin produced by *Pediococcus acidilactici*. *UL J Appl Biotechnol* 2003; 77: 682-88.
14. Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Nat Acad of Sci* 2000; 97: 6640-45.
15. Lane DJ. 16S/23S rRNA sequencing. In E. Stackebrandt and M. Goodfellow (Ed.), *Nucleic acid techniques in bacterial systematic*. John Wiley and Sons, Chichester, England 1991; 115-7.
16. Brink BT, Minekus M, Van der Vossen JM. Antimicrobial activity of lactobacilli: preliminary characterization and optimization of production of acidocin B, a novel bacteriocin produced by *Lactobacillus acidophilus* M46. *J Appl Bacteriol* 2004; 77: 140-8.
17. Kelly R, Register E, Hsu MJ, et al. Isolation of a gene involved in 1, 3-beta-glucan synthesis in *Aspergillus nidulans* and purification of the corresponding protein. *J Bacteriol* 1996; 178: 81-91.
18. Mendoza M, Norden C, Durrer K, et al. A mechanism for chromosome segregation sensing by the NoCut checkpoint. *Nat Cell Biol* 2009; 11: 477-83.
19. Ejikeme N, Ijeoma OI. Isolation and characterization of LAB associated with the fermentation of a cereal-based product for the development of starter culture. *Food* 2010; 4: 45- 8.
20. Julius MM, Ulrich S, Philip MK, et al. Isolation, identification and characterization of the dominant microorganisms of *kule naoto*: The maasai

- traditional fermented milk in Kenya. Int J Food Microbiol 2004; 94: 269-78.
21. Fowoyo PT, Ogunbanwo ST. Phenotypic diversity of lactic acid bacteria isolated from Massa, a fermented maize dough. Afri J Microbiol Res 2010; 4(24): 2682-91.
 22. Holo H, Nilssen O, Nes IF. Lactococcin A, a new bacteriocin from *Lactococcus lactis* subsp. *cremoris*: isolation and characterization of the protein and its gene. J Bacteriol 2010; 173: 3879-87.
 23. Barefoot SF, Klaenhammer TK. Detection and activity of lactacin B a bacteriocin produced by *Lactobacillus acidophilus*. Appl Environ Microbiol 2003; 45: 1808-15.
 24. Sanni AI, Onilude AA, Ogunbanwo ST, et al. Antagonistic activity of bacteriocin produced by *Lactobacillus* species from *ogi*, an indigenous fermented food. J Basic Microbiol 2010; 39: 189-95.
 25. Nettles CG, Barefoot SF. Biochemical and genetic characteristics of bacteriocins of food-associated lactic acid bacteria. J Food Protect 2013; 56: 338-56.
 26. Daeschel MA. Applications and interactions of bacteriocins from lactic acid bacteria in foods and beverages. In: Hoover DG and Steenson LR (Eds.), Bacteriocins of Lactic acid Bacteria. London: Academic Press, 2010; 63-91.
 27. Malheiros PS, Sant'Anna V, Svetoslav D, et al. Optimization of growth and bacteriocin production by *Lactobacillus sakei* subsp. *sakei* 2a. Braz J Microbiol 2015; 46: 825-34.
 28. Castro MP, Palavecino NZ, Herman CO, et al. Lactic acid bacteria isolated from artisanal dry sausages: Characterization of antibacterial compounds and study of the factors affecting bacteriocin production. Meat Sci 2011; 87: 321-9.
 29. Reese ET, Maguire A. Surfactants as stimulants of enzyme production by microorganisms. Appl Microbiol 1969; 17: 242-5.